



Marshall Islands - Outgoing Up To 24 Persons						
Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M1	1. Aspirin 300mg tablets	To relieve aches and pains in diseases such as influenza, the common cold, and sinusitis.	For Headache, 1-3 300-mg tablets, repeated in 6 hours, if necessary. For pains and aches in joints associated with neuralgia, colds and other ailments, 2 tablets administered with a 30-mg tablet of codeine sulphate may be indicated	600		
M2	2. Alcohol, rubbing 70% Isopropyl - 16oz bottles	As an antiseptic for "degerming" the skin prior to injections and other surgical procedures. For disinfecting hands, rub the skin gently but thoroughly for 3 minutes with gauze or cloth soaked with the alcohol.	As directed by Doctor or Medical Officer.	6		
M3	3. Aluminium Acetic Acid 2% Otic solution (Domeboro) 60ml	As Directed	As directed by Doctor or Medical Officer.	2		
M4	4. Aluminium Acetate powder (Domeboro) - 12oz	For external use only. A solution / mixture, prepared from a powder pack as indicated on the carton, is used as a soothing, astringent wet dressing for inflamed or "weeping" areas of the skin. It is an effective, simple remedy for inflammation of areas of the skin due to insect bites, poison ivy, or certain fungal infections such as athletes foot (acute dermatophytosis)	Mix 1 or 2 packets in 500ml of water, as directed on the package for specific conditions, or as directed by a physician. DO NOT strain or filter. the solution is applied in the form of wet compress to the affected areas; this should be repeated several times a day, each time for at least an hour. Also, it may be used as a soak for hands or feet in acute infections such as contact eczema or athletes foot.	2		
M5	5 Alumina and magnesia Gel (Maalox) - 12oz	Used for treating gastric hyperacidity and peptic ulcer, acid indigestion, or sour stomach. Generally promotes healing and relieves pain. Neutralizes hydrochloric acid in the stomach without producing an alkaline reaction. The liquid form of the drug is more effective than the tablet form.	10ml (about 2 teaspoonfuls) 2 hours after meals and at bedtime. Dose and frequency depend on severity of symptoms and relief obtained. The dose range is 5-30ml per administration. In severe gastritis it can be given every 2 hours, possibly with half a glass of milk.	6		
M6	6. Alumina and Magnesia tablets (Maalox) - 100 tablets	Same as for the oral suspension; however, somewhat less effective.	1 or 2 tablets, chewed thoroughly before swallowing, 2-4 hours after meals, as needed, and at bedtime.	1000		
M7	7. Aminophylline 500mg Suppositories Disc By Mfg, - 10 pack	To facilitate breathing in patients who have bronchial asthma, asthmatic bronchitis, pulmonary emphysema, and certain types of heart failure. Wheezing is usually an indication for use. The drug also has diuretic activity (increases urination) give only on RADIO MEDICAL ADVICE	Insert one 500-mg suppository rectally. This dose may be repeated in 8-12 hours. The total dose should not exceed 1g (2 suppositories) in 24 hours.	20		
M8	8. Amitriptyline 25mg Tablets (elavil) 100 tablets	A psychotherapeutic drug used to treatment of depression and anxiety. It is a slow acting drug, and should be used only after obtaining RADIO MEDICAL ADVICE. Amitriptyline has sedative as well as antidepressive properties, so do not give any other sedative along with it,	50mg orally the first day before bedtime, and the same dose morning and evening on the second and consecutive days.	100		
M9	9. Ampicillin capsules, 250mg, 100's	As Directed	As directed by Doctor or Medical Officer.	300		



Marshall Islands - Outgoing Up To 24 Persons						
Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M10	10. Hemorrhoidal Suppositories - 12 pack	For temporary relief of itching, burning and soreness from haemorrhoids (piles). Soothes inflamed haemorrhoids. The patient should try to avoid straining on the stool, heavy lifting coughing, and excessive sneezing. Control constipation (so that straining should not be necessary) with mild laxatives. Warm baths may help.	Remove wrapper and insert one suppository as deeply as possible into the rectum in the morning, at bedtime, and immediately after each bowel movement.	72		
M11	11. Vitamin C 250mg Tablets (Ascorbic Acid) - 100 tablets	Febrile states, chronic illness, and infection may increase the need for this vitamin. Deficiency of this vitamin causes scurvy. Symptoms of mild deficiency include bleeding gums and loosened teeth.	As directed by Doctor or Medical Officer.	300		
M12	12. Atropine sulphate 0.4mg/ml injection - 1ml units	Injectable atropine sulphate is included for use as an antidote in the treatment of poisoning by insecticides containing organophosphate or carbamate chemicals as part of their formulation. get RADIO MEDICAL ADVICE in the event of such poisoning.	As directed by Doctor or Medical Officer.	60		
M13	13. Ampicillin 1 gm injection - each	As Directed	As directed by Doctor or Medical Officer.	20		
M14	14. Tolnaftate 1% cream (Tinactin) - 15gm tube	As Directed	As directed by Doctor or Medical Officer.	4		
M15	15. Calamine lotion - 4oz size	For relieving itching or irritated skin, heat rash, and hives. Shake the bottle well. Wet a pad of cotton with the lotion and pat it on the affected area.	As directed by Doctor or Medical Officer.	8		
M16	16. Calcium Gluconate 648mg Tablets each	This medicament is intended to be used primarily in cases of poisoning by oxalic acid or sodium fluoride. This sort of poisoning may cause a depletion of the calcium content of the blood, resulting in convulsions. Calcium gluconate acts by restoring needed calcium to the blood.	Dissolve 5 tablets in 250ml of water, to be drunk immediately.	30		
M17	17. Hibiclens Solution (Chlorhexidine Gluconate) - 16oz	As Directed	As directed by Doctor or Medical Officer.	3		
M18	18. Charcoal Activated Powder 227g	In the initial treatment of most poisonings (because of its property to absorb many poisons)	10g (2 tablespoonfuls). Mix the powder with water prior to administration. Re-peat if the patient vomits.	1		
M19	19. Chloroquine 250mg Tablets - 100 tablets	To prevent and treat malaria. When a ship is in a known malarial area, obtain medical advice from a physician on the prophylactic measures effective in that region	As directed by Doctor or Medical Officer.	100		
M20	20. Chlorphenamine Maleate 4mg Tablets - 100 tablets	For treating allergic reactions, such as hayfever, urticaria, and anaphylactic shock.	1 tablet, one to three times daily.	100		
M21	21. Diphenhydramine 50mg/ml injection (Benadryl) 1ml units	As Directed	As directed by Doctor or Medical Officer.	20		
M22	22. Chlorpromazine 25mg Tablets (Thorazine) - each	For treating anxiety, tension, and agitation: also to control nausea and vomiting.	25mg for the control of vomiting for cases of obvious madness, give 1 tablet three times a day and consult the doctor, who may advise increaing the dosage	100		
M23	23. Chlorpromazine 25mg/ml injection (Thorazine) - 1ml units	Same as chlorpromazine tablets	50mg (2 ampoules) should be given to patients with delirium tremens this dose may be repeated after 6 hours. in ther cases 1 or 2 25 mg njections daily	20		



Marshall Islands - Outgoing Up To 24 Persons						
Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M24	24. Clove oil - 1oz	For the relief of toothache when there is a cavity in the tooth. If possible, dry the cavity with cotton wrapped on a toothpick and then pack it with a piece of cotton that has been dipped in oil of cloves. This procedure may be repeated as often as necessary. Because clove oil has an irritating effect on tissues, use extra care to avoid contact with surrounding gum or other mouth areas	As directed by Doctor or Medical Officer.	2		
M25	25. Acetaminophen W/Codeine 30mg tablets (tylenol) - 100 tablets	To relieve coughing (2) in diarrhoea	For the persistent and severe coughing accompanying severe respiratory infections, give half a 30-mg tablet as often as every 2 hours, if necessary. This time interval should be lengthened as soon as the cough is controlled. Codeine sulfate should be discontinued as soon as the cough is relieved. In diarrhoea, 30mg may be given, repeated after 4 hours if necessary.	100		
M26	26. Meclizine 25mg tablets (Antivert) - 100 tablets	As Directed	As directed by Doctor or Medical Officer.	400		
M27	27. Dextran 6% & 0.9% Sodium Chloride IV Injection - 500ml units	To expand plasma volume and maintain blood pressure in the emergency treatment of shock caused by a loss of blood. Also used in the treatment of burns. Dextran should be purchased in a package that includes an administration set.	Give only on RADIO MEDICAL ADVICE from a physician. 500ml infused intravenously at the rate of 20ml per minute, so that the entire amount is given over a period of about 30 minutes. The total amount administered should not exceed 20ml per Kg of body weight during the first 24 hours.	6		
M28	28. Diazepam 5mg/ml Injection (Valium) - 2ml units	(1) For treatment of severe agitation including acute alcohol withdrawal states and convulsions (epilepsy) (2) May be useful for shivering due to generalized hypothermia.	The usual dosage is 2-10 mg (intramuscularly, deep into the muscle), Which may be repeated once in 1-4 hours, depending upon the response and the severity of the condition.	20		
M29	29. Diazepam 5mg tablets(Valium) - 100 tablets	(1) For treatment of common anxiety and tension , and (2) For the management of agitation during alcohol withdrawal.	Varies from 2 to 10mg 2-4 times a day, depending upon the condition being treated.	200		
M30	30. Dimercaprol 100mg/ml Injection - 2ml units	The drug is used as an antidote in poisoning. Indicated in the treatment of arsenic (pesticides), gold and mercury poisonings also in acute lead poisoning, in combination with calcium disodium edetate. Always obtain RADIO MEDICAL ADVICE in the event of such poisoning.	As directed by Doctor or Medical Officer.	6		
M31	31. Doxycycline 100mg Capsules - 50 capsules	Antibacterial agent used in the treatment of urinary infections. It is given in cases of both gonococcal and nongonococcal urethritis after spectinomycin hydrochloride or sulfamethoxazole / trimethoprim has been used.	The usual dose for adults is 200mg on the first day of treatment, given as 100mg every 12hours, followed by a maintenance dose of 100mg daily. In severe infections e.g. chronic infection of the urinary tract, 100mg every 12 hours is recommended	200		



Marshall Islands - Outgoing Up To 24 Persons

Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M32	32. Ephedrine Sulfate 25mg Capsules - 100 capsules	(1) To relieve difficult breathing in asthma (2) to prevent asthmatic attacks in chronic cases (effects appear 30-60 minutes after administration) and (3) to relieve nasal congestion in hayfever and severe head colds. obtain RADIO MEDICAL ADVICE before using this drug	One 25mg capsule four times a day. If ephedrine is used for several days, phenobarbital may be indicated to overcome its stimulant effects (one 30mg phenobarbital tablet up to three times a day)	100		
M33	33. Epinephrine 1mg/ml injection - 1ml units	(1) For acute asthma attacks and for severe allergic reactions to penicillin and other drugs, and to insect bites (2) For cardiac or circulatory failure	By subcutaneous or intramuscular injection, 0.3 ml of 1:1000 solution, every 2 hours as necessary. The usual dosage range is 0.1-0.5 ml in asthma, and up to 1ml in other conditions	20		
M34	34. Methergine 0.2 mg/1ml injection - 1ml units	As Directed	As directed by Doctor or Medical Officer.	10		
M35	35. Erythromycin 250mg tablets - 100 tablets	For a variety of infections of the upper and lower respiratory tract; infections of the mouth, gums, and teeth; infections of the nose, ears, and sinuses. Give this antibiotic only when RADIO MEDICAL ADVICE has been received from a physician, and for some infections resistant to penicillin.	One 250-mg tablet, four times a day (for serious infections, dosage may be increased to two 250mg tablets, four times a day)- to be continued for 48 hours or more after symptoms have subsided.	300		
M36	36. Tetracaine 0.5% Ophth Solution (Pontocaine) - 15ml units	For local anaesthesia of the eye, before removing foreign bodies.	Put 3 drops inside the eyelids, repeat this three times at 2 minute intervals.	2		
M37	37. Triple Antibiotic Pphth Solution - 10ml units	As Directed	As directed by Doctor or Medical Officer.	3		
M38	38. Triple Antibiotic Ophth ointment (Neosporin) 3.5gm	As Directed	As directed by Doctor or Medical Officer.	6		
M39	39. Eye wash Sterile -4oz	For irrigating or flushing the eye to wash away foreign particles, mucous secretions, and fluorescein dye used in diagnosis. Before use, warm the bottle with the solution to body temperature.	As directed by Doctor or Medical Officer.	6		
M40	40. Fluorets Sodium Eye Strips 100 per box	As Directed	As Directed	#		
M41	41. Furosemide tablets, 40mg, (Lasix) 100's.	Furosemide is a potent short-acting diuretic, which causes the production of urine by affecting the kidneys. It is indicated in the treatment of excess body fluid (oedema) associated with congestive heart failure, cirrhosis of the liver, or certain kidney diseases.	The usual oral dosage for adults is 20-80 mg in a single dose in the morning, for 2-4 consecutive days each week, followed by a drug-free period. However, the dosage will vary, depending on the disease and the patients response.	100		
M42	42. Nitro-Quick 0.4mg Sublingual Tablets - 25	As Directed	As directed by Doctor or Medical Officer.	50		
M43	43. Solu-Cortef 100mg Vial Injection (Hydrocortisone) - each	As Directed	As directed by Doctor or Medical Officer.	5		
M44	44. Hydrocortisone 1% Ointment - 1oz	For temporary relief of certain skin disorders, common rashes, inflamed skin and disorders causing itching and discomfort. It may be used for temporary relief of itching, burning and soreness from haemorrhoids (piles).	Apply a thin film to the affected skin 2-4 times a day. Apply sparingly and with gentle rubbing. Clean the skin before each use	6		
M45	45. Ichtammol 10% Ointment - 1oz	As Directed	As directed by Doctor or Medical Officer.	6		
M46	46. Insect repellent pump - 2oz	External use only. To repel mosquitos, chiggers, flies, and other biting insects. It can be used on clothing or the skin, protecting for up to 8 hours.	As directed by Doctor or Medical Officer.	12		
M47	47. Iodine Tincture 2% Mild - 1oz	As Directed	As directed by Doctor or Medical Officer.	10		
M48	Xylocaine 1% 10mg/ml Injection (Lidocaine) - 2ml units	As Directed	As directed by Doctor or Medical Officer.	12		
M49	49. Lindane lotion (Kwell) - 2oz	As Directed	As directed by Doctor or Medical Officer.	12		



Marshall Islands - Outgoing Up To 24 Persons						
Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M50	50. Milk of Magnesia - 12oz	As Directed	As directed by Doctor or Medical Officer.	8		
M51	51. Metronidazole 250mg tablets (Flagyl) - 100 tablets	(1) For treatment of salpingitis and other diseases of women and (2) for the treatment of amoebic dysentery.	(1) for diseases of women, usually 1 tablet every 8 hours for 2 weeks. (2) For the dosage in amoebic dysentery, get RADIO MEDICAL ADVICE	500		
M52	52. Miconazole Vaginal Cream (micatin) - 45gm	As Directed	As directed by Doctor or Medical Officer.	5		
M53	53. Miconazole Vaginal Suppositories 7 W/Applicator - each	As Directed	As directed by Doctor or Medical Officer.	3		
M54	54. Mineral oil Heavy - 16oz	Mineral oil (liquid petrolatum) is used as a skin emollient or protective agent. It may also be used to help remove ointments or creams from treated skin areas. Its administration by mouth may be indicated after the ingestion of certain poisons, including alkalis.	Apply liberally over areas of the skin.	1		
M55	55. Morphine sulphate injection, 10mg/ml, 1-ml units	For severe pain not relieved by other analgesics. OBTAIN RADIO MEDICAL ADVICE prior to use. Discontinue as soon as the pain can be relieved by other drugs that can be given orally and are less addictive.	For the relief of severe pain following injuries or burns and severe pain of sudden origin in the abdomen or chest, give 10-15mg intramuscularly. If the pain is unrelieved, or if it recurs soon after the first dose, a second dose of 10 mg intramuscularly, may be given 1 hour or more later. Give further doses if necessary, at intervals of at least 4 hours not sooner.	20		
M56	56. Naloxone Hydrochloride 0.4 mg/ml Injection (Narcan) - 1ml units	For the emergency treatment of respiratory depression resulting from the administration of narcotics, such as morphine. Also indicated for the detection of suspected acute narcotic over-dose.	May be given by subcutaneous, intramuscular, or intravenous injection. The intravenous route of administration is recommended only for emergency situations. the usual dose is 0.4mg, repeated every 2 or 3 minutes until a favourable response is achieved. If no improvement is noted after 2 or 3 doses, another cause of the depression should be suspected.	6		
M57	57. Triple Antibiotic Ointment (Neosporin) - 1oz	As Directed	As directed by Doctor or Medical Officer.	20		
M58	Electrolyte Tablets - 100 tablets	As Directed	As directed by Doctor or Medical Officer.	100		
M59	59. Oxygen Refil Size "D" each	As Directed	As directed by Doctor or Medical Officer.	2		
M60	60. Acetaminophen 500mg tablets (tylenol) - 100 tablets	As Directed	As directed by Doctor or Medical Officer.	300		
M61	61. Petrolatum Ointment - 1oz	As a bland and neutral protective dressing. Apply to minor burns, or dry skin.	As directed by Doctor or Medical Officer.	12		



Marshall Islands - Outgoing Up To 24 Persons						
Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M62	62. Phenobarbital 30mg Tablets (Tylenol) - 100 tablets	(1) For insomnia (2) for relatively prolonged sedation, as in some anxiety and tension states and in hypertension and (3) for treatment of epilepsy.	To induce sleep, 90mg at bedtime. As a sedative, one 30mg tablet, 1-4 times daily. In epilepsy, the dose should be adjusted to the individual. consult the doctor in each case. The average range for an adult is 50-120mg daily, although much higher doses may be required for short periods of time	300		
M63	63. Penicillin VK 250mg Tablets - 100 tablets	As Directed	As directed by Doctor or Medical Officer.	300		
M64	64. Pilocarpine 2% Opth Solution (Isopto Carpine) - 15ml units	For the management of glaucoma. Pilocarpine hydrochloride eye-drops constrict the pupil. It is anticipated that this medicament will be needed aboard ship only for patients who have run out of their own supply.	For glaucoma, 2 drops in the eye every 6 hours daily, or as prescribed	2		
M65	65. Potassium Permanganate Crystal - 500g	Diluted un water (1g of crystals to 0.5-1 litre of water.) This substance is used for the disinfection of the skin and mucous membrabes and for the treatment of some skin diseases.	(1g of crystals to 0.5-1 litre of water	1		
M66	66. Probenecid 500mg Tablets (Benemid) - 100 tablets	As an adjunct to intensive therapy with benzylpenicillin, with ampicillin, and a number of other penicillins, for prolonging the effective blood concentration of these drugs	Generally one 500mg tablet four times a day, as long as penicillin is taken.	200		
M67	67. Ampicillin 500mg Injection - each	As Directed	As directed by Doctor or Medical Officer.	20		
M68	68. Proguanil 100mg (Pauludrine) - 100	For prevention of malaria, given either alone or in association with chloroquine. When a ship is in known malarial areas, obtain medical advice on the prophylactic measures effective in that areas.	As directed by Doctor or Medical Officer.	100		
M69	69. Pinworm Caplets (Pyrantel) - 24	For the treatment of (1) pinworm infections and (2) roundworm infections in adults and children	Orally, in a single dose of 10mg per kg of body weight (2 tablets fro a patient weighing 50kg). The highest single dose should not exceed 4 tablets (1g). May be administered with out regard to ingestion of food or time of day. May be taken with either milk or fruit juice	48		
M70	70. Quinine Sulfate 325mg capsules - 100 capsules	In the treatment of acute attacks of malaria contracted in certain areas. Quinine sulfate is not administered to prevent or suppress infection.	For acute malaria attacks, give 2 tablets every 8 hours, until the patient sees a doctor. If the treatment has to be continued for more than 10 days, get RADIO MEDICAL ADVICE. This medicine should be taken after meals to lessen the possibility of stomache upsets.	200		
M71	71. Quinine Dihydrochloride Injection 300mg/ml 2ml units	In the treatment of acute attacks of malaria when the patient is unable to tolerate oral medication because of repeated vomiting or impairment of consciousness.	For acute malaria attacks in adult patients unable to take oral drugs, give 600mg intramuscularly (in the upper outer quadrant of the buttock) every 8 hours until the patient can take oral medication. At that time stop the injections and give quinine sulfate tablets.	20		



Marshall Islands - Outgoing Up To 24 Persons						
Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M72	72. Albuterol 17g Inhalation (Proventil) - 200 metered doses	As Directed	As directed by Doctor or Medical Officer.	2		
M73	73. Sodium Chloride 0.9% IV Injection - 1000ml units	Intravenously (1) to replace water, chloride, and sodium lost in extensive vomiting or diarrhoea, (2) in dehydration due to excessive sweating and (3) in mild haemorrhage. OBTAIN RADIO MEDICAL ADVICE before administering.	Depends on patients condition. Generally, 1500-3000 ml may be given over 24 hours. Do not give intravenous solutions at a faster rate than 500ml per hour, except on RADIO MEDICAL ADVICE	6		
M74	74. IV Solution set vented W/Injection site 15 drop/ml each	As Directed	As directed by Doctor or Medical Officer.	6		
M75	75. IV Solution set 15 Drops/ml Macro drip - each	As Directed	As directed by Doctor or Medical Officer.	6		
M76	76. Thermotabs (enteric coated salt tablets) - 100 tablets	As Directed	As directed by Doctor or Medical Officer.	200		
M77	77. Trobicin Sterile Powder Injection (Spectinomycin)	As Directed	As directed by Doctor or Medical Officer.	20		
M78	78. Sulfamethoxazole/Trimethoprim 400mg/80mg tabs (Bactrim) - 100 tablets	For the treatment of urinary infections, after obtaining RADIO MEDICAL ADVICE	1-2 tablets every 12 hours, for 7 days. Give each dose with full glass of water.	200		
M79	79. Baby Powder J & J (Talc) - 4oz	As as skin lubricant and protector, and as a dusting powder for such skin irritations as chafing and prickly heat	As directed by Doctor or Medical Officer.	6		
M80	80. Tetanus immune Globulin (human) Prefilled syringe - 250 units	To provide passive immunity to tetanus in a patient who is not protected through vaccination with adsorbed tetanus toxoid and has sustained a potentially contaminated wound.	For passive immunization, 250 units by intramuscular injection only.	5		
M81	81. Tetanus Toxoid Prefilled syringe Injection - 0.5ml units	For active immunization against tetanus	For primary immunization of adults and children over 6 years of age. 3 doses are required (see package insert for amount of each dose-usually 0.5 or 10.0ml). The toxoid is injected intramuscularly into the lateral aspect of the upper arm. The second dose is given 4-6 weeks after the first, and the third dose is 6 months to 1 year after the second thereafter, the recommended dose is administered every 5 years.	10		
M82	82. Antibiotic Otic Solution (Generic Cortisporin) - 10ml units	As Directed	As directed by Doctor or Medical Officer.	10		
M83	83. Tetracycline 250mg capsules (Achromycin) - 100 capsules	For susceptible infections; to be given after receiving RADIO MEDICAL ADVICE. do not give to children under 12 years of age.	Usually, one 250mg capsule every 6 hours. In severe illness, increase to two 250-mg capsules every 6 hours.	300		
M84	84. Water for injection Sterile - 5ml units	To dissolve penicillin in powder form	As directed by Doctor or Medical Officer.	30		
M85	85. Zinc oxide ointment	A nontoxic, protective, water-insoluble mildly astringent, and antiseptic paste employed in a large variety of diseases and irritations of the skin.	As directed by Doctor or Medical Officer.	12		
M86	Emesis Basin - Each	As Directed	As Directed	1		
M87	87. Wash Basin Plastic - Large	As Directed	As Directed	1		
M88	88. Bed pan Fracture - each	As Directed	As Directed	1		
M89	89. Cane Aluminum Adjustable #A756 each	As Directed	As Directed	1		
M90	90. Crutches Aluminum Complete - adult regular	As Directed	As Directed	1		
M91	91. Eye Cup Glass each	As Directed	As Directed	2		
M92	92. Funnel Stainless steel - 6oz	As Directed	As Directed	1		
M93	93. Cylinder Glass Double Scale - 50ml	As Directed	As Directed	1		



Marshall Islands - Outgoing Up To 24 Persons

Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M95	95. Heating Pad Cara Dry #50 - each	As Directed	As Directed	1		
M96	96. Hot water bottle / icebag 2 quart each	As Directed	As Directed	1		
M97	97. Strokes litter basket - each	As Directed	As Directed	1		
M98	98. Magnifying glass 2-1/2" - each	As Directed	As Directed	1		
M99	99. Stretcher neil robertson - each	As Directed	As Directed	1		
M100	100. Oxygen unit (complete) W/Case fro "D" size tank - each	As Directed	As Directed	1		
M101	101. Oxygen Cylinder Size "D" empty - each	As Directed	As Directed	1		
M102	102. Refrigerator Walnut colour - Each	As Directed	As Directed	1		
M103	103. Restraints Wrist/Ankle padded pair - each	As Directed	As Directed	1		
M104	104. Straight Jacket	As Directed	As Directed	1		
M105	105. Bedrails Home style #32-GF6408 - 1pair	As Directed	As Directed	1		
M106	106. Resuscitator bag adult disp. w/Mask & tubing	As Directed	As Directed	1		
M107	107. Inflatable ring rubber	As Directed	As Directed	1		
M108	108. Scale doctor plus 0-300lb	As Directed	As Directed	1		
M109	109. Sphygmomanometer aneroid #115	As Directed	As Directed	1		
M110	110. Splint inflatable kit - 4 per kit	As Directed	As Directed	1		
M111	111. Finger Splint padded assorted sizes	As Directed	As Directed	2		
M112	112. Sterilizer Dri-Clave 15-1/2x10-1/2x9-1/2 115 volt	As Directed	As Directed	1		
M113	113. Stethoscope Black - 22"	As Directed	As Directed	1		
M114	114. Thermometer Bath flat wood #50-1537	As Directed	As Directed	2		
M115	115. Instrument Tray Stainless steel - 8-1/2 x 3x1-1/2	As Directed	As Directed	1		
M116	116. Urinal male disposable	As Directed	As Directed	2		
M117	117. Waste recepticle step on 32 Qt white #58-8286	As Directed	As Directed	1		
M118	118. International medical guide for ships	As Directed	As Directed	1		
M119	119. Medical first aid/dangerous goods	As Directed	As Directed	1		
M120	120. International Health regulations	As Directed	As Directed	1		
M121	121. Vaccination Certificate Manual (WHO)	As Directed	As Directed	1		
M122	122. Airway Kit Nasopharyngeal - 5 sizes w/case	As Directed	As Directed	1		
M123	123. Scalpel Blades #10	As Directed	As Directed	6		
M124	124. Scalpel Blades #15	As Directed	As Directed	6		
M125	125. Forceps Dressing Bayonet - shaped 7"	As Directed	As Directed	1		
M126	126. Forceps Halsted Mosquito Curved #20 - 2673	As Directed	As Directed	2		
M127	127. Forceps Kelly Straight - 5-1/2"	As Directed	As Directed	2		
M128	128. Forceps Kelly Curved -5-1/2"	As Directed	As Directed	2		
M129	129. Forceps Splinter -3-1/2"	As Directed	As Directed	1		
M130	130. Forceps Tissue 1x2 teeth -4-1/2"	As Directed	As Directed	1		
M131	131. Scalpel Handle #3	As Directed	As Directed	2		
M132	132. Scissors Bandage -7-1/2"	As Directed	As Directed	3		
M133	133. Scissors Operating Straight Sharp/Sharp-5-1/2"	As Directed	As Directed	2		
M134	134. Hammer Taylor Type Reflex #15-1305-1	As Directed	As Directed	1		
M135	135. Needle Holder Mayo-Hager 5-1/2"	As Directed	As Directed	2		
M136	136. Tape Micropore Paper 2" x 10yd	As Directed	As Directed	4		
M137	137. Tape Micropore Paper 1" x 10 yd	As Directed	As Directed	4		



Marshall Islands - Outgoing Up To 24 Persons

Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M138	138. IV Butterfly Needle W/Tubing 19g x 3/4"-12"	As Directed	As Directed	6		
M139.	139. Coton Tipped Applicators 6	As Directed	As Directed	200		
M140	140. Elastic Bandage 4" x 4.5 yd	As Directed	As Directed	12		
M141	141. Elastic Bandage 3" x 4.5 yd	As Directed	As Directed	12		
M142	142. Elastic Bandage 2" x 4.5 yd	As Directed	As Directed	12		
M143	143. Elastic Bandage Cotton 2	As Directed	As Directed	12		
M144	144. Flexilite Conforming Gauze Bandage 2" x 4-1/2"	As Directed	As Directed	200		
M145	145. Flexilite Conforming Gauze Bandage 6" x4-1/2"	As Directed	As Directed	40		
M146	146. Flexilite Conforming Gauze Bandage 4" x4-1/2"	As Directed	As Directed	60		
M147	147. Gauze Telfa	As Directed	As Directed	3000		
M148	148. Band Aid Adhesive Surgical Dressing 8" x 6"	As Directed	As Directed	20		
M149	149. Triangular Bandage	As Directed	As Directed	4		
M150	150. Surgitube #2 7/8" x 5 yd	As Directed	As Directed	10		
M151	151. Bandage Spray - 3oz	As Directed	As Directed	4		
M152	152. Vaseline Dressing 3" x 18"	As Directed	As Directed	5		
M153	153. Vaseline Dressing 3" x9"	As Directed	As Directed	10		
M154	154. Vaseline Dressing 6" x 36"	As Directed	As Directed	5		
M155	155. Bandage Compress 4" (1 per box)	As Directed	As Directed	20		
M156	156. Bandage Compress 2" (4 per box)	As Directed	As Directed	3		
M157	157. Bandage Compress 3" (2 per box)	As Directed	As Directed	10		
M158	158. Bottle Plastic - 2oz	As Directed	As Directed	50		
M159	159. Catheter Urethral - 12Fr	As Directed	As Directed	1		
M160	160. Catheter Urethral - 14 Fr	As Directed	As Directed	1		
M161	161. Catheter Urethral - 6 Fr	As Directed	As Directed	1		
M162	162. Catheter Urethral - 8 Fr	As Directed	As Directed	1		
M163	163. Catheter Urethral - 16 Fr	As Directed	As Directed	1		
M164	164. Catheter Tray Urethral - 14 Fr	As Directed	As Directed	3		
M165	165. Clinical Record Chart	As Directed	As Directed	50		
M166	166. Temperature Chart	As Directed	As Directed	50		
M167	167. Medical Report For Seafarers	As Directed	As Directed	100		
M168	168. Cervical Collar Universal - 3"	As Directed	As Directed	1		
M169	169. Cotton Rolled Sterile - 2oz	As Directed	As Directed	3		
M170	170. Cotton Rolled Sterile - 1/2oz	As Directed	As Directed	20		
M171	171. Cotton Rolled Sterile -4oz	As Directed	As Directed	10		
M172	172. Finger Cots Assorted Sizes Sm.,Med., & Large - 12	As Directed	As Directed	12		
M173	173. Penlight Heavy Duty w/batteries	As Directed	As Directed	1		
M174	174. Surgitube #1 5/8" x 5yd	As Directed	As Directed	5		
M175	175. Gloves Latex Surgical "sterile" size 8	As Directed	As Directed	12		
M176	176. Gown Surgical Disposable	As Directed	As Directed	5		
M177	177. Face mask w/elastic band	As Directed	As Directed	50		
M178	178. Medicine Cups plastic 1oz	As Directed	As Directed	100		
M179	179. Microscope slides 1x3 plain	As Directed	As Directed	100		
M180	180. Body bag	As Directed	As Directed	2		
M181	181. Needle Disposable 25g x 5/8"	As Directed	As Directed	36		
M182	182. Needle Disposable 18g x 1-1/2"	As Directed	As Directed	18		
M183	183. Needle Disposable 21g x 1-1/2"	As Directed	As Directed	12		
M184	184. Hematest Reagent tablets	As Directed	As Directed	100		
M185	185. Surgipad Combine Dressing 8"x 10" sterile	As Directed	As Directed	6		
M186	186. Eye pad large sterile - 12 per box	As Directed	As Directed	12		
M187	187. Gauze pads non-adherent 3"x4" sterile - 100 per box	As Directed	As Directed	100		
M188	188. Safety pins Assorted sizes - 50 per	As Directed	As Directed	100		
M189	189. Brush (Surgeons scrub)	As Directed	As Directed	2		
M190	190. Condoms Lubricated	As Directed	As Directed	50		

**Marshall Islands - Outgoing Up To 24 Persons**

Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M191	191. Sheet waterproof 36x72"	As Directed	As Directed	2		
M192	192. Butterfly Closure Medium - 100 each	As Directed	As Directed	1		
M193	193. Steri Strip 1/2"x4" - 6 per	As Directed	As Directed	12		
M194	194. Sutures Chromie Gut 2-0 27" FS1 Needle #637G	As Directed	As Directed	12		
M195	195. Sutures Silk 3-0 18" FS-2 Needle #684G	As Directed	As Directed	24		
M196	196. Sutures Chromic Gut 3-0 27" FS2 Needle #636G	As Directed	As Directed	12		
M197	197. Sutures Nylon 5-0 18" FS-2 Needle #661G	As Directed	As Directed	12		
M198	198. Suture Removal Kit	As Directed	As Directed	1		
M199	199. Glass Syringe 2 ml	As Directed	As Directed	4		
M200	200. Glass Syringe 20ml	As Directed	As Directed	2		
M201	201. Glass Syringe 5 ml	As Directed	As Directed	4		
M202	202. Syringe & Needle 3 cc 21g x 1-1/2"	As Directed	As Directed	50		
M203	203. Syringe & Needle 5 cc 21g x 1-1/2"	As Directed	As Directed	30		
M204	204. Syringe & Needle 3 cc 25g x 5/8"	As Directed	As Directed	50		
M205	205. Chemstrip 9 -100	As Directed	As Directed	100		
M206	206. Clinitest Tablets/Urine Sugar - 36 tablets per	As Directed	As Directed	108		
M207	207. Kleenex - 250 sheets per box	As Directed	As Directed	6		
M208	208. Thermometer Dual scale Oral	As Directed	As Directed	6		
M209	209. Tourniquet Grafkette adult size	As Directed	As Directed	1		
M210	210. Tongue Depressors Wood 6" Senior	As Directed	As Directed	100		
M211	211. Towels Poly-lined 18"x26"	As Directed	As Directed	24		
M212	212. Towels Cotton 18"x33" - 12 each	As Directed	As Directed	12		
M213	213. Sterilized pouches tower self seal 3.5x8" - 100 per	As Directed	As Directed	100		
M214	214. "Sharps" disposal box	As Directed	As Directed	1		



Marshall Islands - Outgoing Up To 24 Persons						
Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M1	1. Aspirin 300mg tablets	To relieve aches and pains in diseases such as influenza, the common cold, and sinusitis.	For Headache, 1-3 300-mg tablets, repeated in 6 hours, if necessary. For pains and aches in joints associated with neuralgia, colds and other ailments, 2 tablets administered with a 30-mg tablet of codeine sulphate may be indicated	600		
M2	2. Alcohol, rubbing 70% Isopropyl - 16oz bottles	As an antiseptic for "degerming" the skin prior to injections and other surgical procedures. For disinfecting hands, rub the skin gently but thoroughly for 3 minutes with gauze or cloth soaked with the alcohol.	As directed by Doctor or Medical Officer.	6		
M3	3. Aluminium Acetic Acid 2% Otic solution (Domeboro) 60ml	As Directed	As directed by Doctor or Medical Officer.	2		
M4	4. Aluminium Acetate powder (Domeboro) - 12oz	For external use only. A solution / mixture, prepared from a powder pack as indicated on the carton, is used as a soothing, astringent wet dressing for inflamed or "weeping" areas of the skin. It is an effective, simple remedy for inflammation of areas of the skin due to insect bites, poison ivy, or certain fungal infections such as athletes foot (acute dermatophytosis)	Mix 1 or 2 packets in 500ml of water, as directed on the package for specific conditions, or as directed by a physician. DO NOT strain or filter. the solution is applied in the form of wet compress to the affected areas; this should be repeated several times a day, each time for at least an hour. Also, it may be used as a soak for hands or feet in acute infections such as contact eczema or athletes foot.	2		
M5	5 Alumina and magnesia Gel (Maalox) - 12oz	Used for treating gastric hyperacidity and peptic ulcer, acid indigestion, or sour stomach. Generally promotes healing and relieves pain. Neutralizes hydrochloric acid in the stomach without producing an alkaline reaction. The liquid form of the drug is more effective than the tablet form.	10ml (about 2 teaspoonfuls) 2 hours after meals and at bedtime. Dose and frequency depend on severity of symptoms and relief obtained. The dose range is 5-30ml per administration. In severe gastritis it can be given every 2 hours, possibly with half a glass of milk.	6		
M6	6. Alumina and Magnesia tablets (Maalox) - 100 tablets	Same as for the oral suspension; however, somewhat less effective.	1 or 2 tablets, chewed thoroughly before swallowing, 2-4 hours after meals, as needed, and at bedtime.	1000		
M7	7. Aminophylline 500mg Suppositories Disc By Mfg, - 10 pack	To facilitate breathing in patients who have bronchial asthma, asthmatic bronchitis, pulmonary emphysema, and certain types of heart failure. Wheezing is usually an indication for use. The drug also has diuretic activity (increases urination) give only on RADIO MEDICAL ADVICE	Insert one 500-mg suppository rectally. This dose may be repeated in 8-12 hours. The total dose should not exceed 1g (2 suppositories) in 24 hours.	20		
M8	8. Amitriptyline 25mg Tablets (elavil) 100 tablets	A psychotherapeutic drug used to treatment of depression and anxiety. It is a slow acting drug, and should be used only after obtaining RADIO MEDICAL ADVICE. Amitriptyline has sedative as well as antidepressive properties, so do not give any other sedative along with it,	50mg orally the first day before bedtime, and the same dose morning and evening on the second and consecutive days.	100		
M9	9. Ampicillin capsules, 250mg, 100's	As Directed	As directed by Doctor or Medical Officer.	300		



Marshall Islands - Outgoing Up To 24 Persons						
Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M10	10. Hemorrhoidal Suppositories - 12 pack	For temporary relief of itching, burning and soreness from haemorrhoids (piles). Soothes inflamed haemorrhoids. The patient should try to avoid straining on the stool, heavy lifting coughing, and excessive sneezing. Control constipation (so that straining should not be necessary) with mild laxatives. Warm baths may help.	Remove wrapper and insert one suppository as deeply as possible into the rectum in the morning, at bedtime, and immediately after each bowel movement.	72		
M11	11. Vitamin C 250mg Tablets (Ascorbic Acid) - 100 tablets	Febrile states, chronic illness, and infection may increase the need for this vitamin. Deficiency of this vitamin causes scurvy. Symptoms of mild deficiency include bleeding gums and loosened teeth.	As directed by Doctor or Medical Officer.	300		
M12	12. Atropine sulphate 0.4mg/ml injection - 1ml units	Injectable atropine sulphate is included for use as an antidote in the treatment of poisoning by insecticides containing organophosphate or carbamate chemicals as part of their formulation. get RADIO MEDICAL ADVICE in the event of such poisoning.	As directed by Doctor or Medical Officer.	60		
M13	13. Ampicillin 1 gm injection - each	As Directed	As directed by Doctor or Medical Officer.	20		
M14	14. Tolnaftate 1% cream (Tinactin) - 15gm tube	As Directed	As directed by Doctor or Medical Officer.	4		
M15	15. Calamine lotion - 4oz size	For relieving itching or irritated skin, heat rash, and hives. Shake the bottle well. Wet a pad of cotton with the lotion and pat it on the affected area.	As directed by Doctor or Medical Officer.	8		
M16	16. Calcium Gluconate 648mg Tablets each	This medicament is intended to be used primarily in cases of poisoning by oxalic acid or sodium fluoride. This sort of poisoning may cause a depletion of the calcium content of the blood, resulting in convulsions. Calcium gluconate acts by restoring needed calcium to the blood.	Dissolve 5 tablets in 250ml of water, to be drunk immediately.	30		
M17	17. Hibiclens Solution (Chlorhexidine Gluconate) - 16oz	As Directed	As directed by Doctor or Medical Officer.	3		
M18	18. Charcoal Activated Powder 227g	In the initial treatment of most poisonings (because of its property to absorb many poisons)	10g (2 tablespoonfuls). Mix the powder with water prior to administration. Re-peat if the patient vomits.	1		
M19	19. Chloroquine 250mg Tablets - 100 tablets	To prevent and treat malaria. When a ship is in a known malarial area, obtain medical advice from a physician on the prophylactic measures effective in that region	As directed by Doctor or Medical Officer.	100		
M20	20. Chlorphenamine Maleate 4mg Tablets - 100 tablets	For treating allergic reactions, such as hayfever, urticaria, and anaphylactic shock.	1 tablet, one to three times daily.	100		
M21	21. Diphenhydramine 50mg/ml injection (Benadryl) 1ml units	As Directed	As directed by Doctor or Medical Officer.	20		
M22	22. Chlorpromazine 25mg Tablets (Thorazine) - each	For treating anxiety, tension, and agitation: also to control nausea and vomiting.	25mg for the control of vomiting for cases of obvious madness, give 1 tablet three times a day and consult the doctor, who may advise increaing the dosage	100		
M23	23. Chlorpromazine 25mg/ml injection (Thorazine) - 1ml units	Same as chlorpromazine tablets	50mg (2 ampoules) should be given to patients with delirium tremens this dose may be repeated after 6 hours. in ther cases 1 or 2 25 mg njections daily	20		



Marshall Islands - Outgoing Up To 24 Persons						
Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M24	24. Clove oil - 1oz	For the relief of toothache when there is a cavity in the tooth. If possible, dry the cavity with cotton wrapped on a toothpick and then pack it with a piece of cotton that has been dipped in oil of cloves. This procedure may be repeated as often as necessary. Because clove oil has an irritating effect on tissues, use extra care to avoid contact with surrounding gum or other mouth areas	As directed by Doctor or Medical Officer.	2		
M25	25. Acetaminophen W/Codeine 30mg tablets (tylenol) - 100 tablets	To relieve coughing (2) in diarrhoea	For the persistent and severe coughing accompanying severe respiratory infections, give half a 30-mg tablet as often as every 2 hours, if necessary. This time interval should be lengthened as soon as the cough is controlled. Codeine sulfate should be discontinued as soon as the cough is relieved. In diarrhoea, 30mg may be given, repeated after 4 hours if necessary.	100		
M26	26. Meclizine 25mg tablets (Antivert) - 100 tablets	As Directed	As directed by Doctor or Medical Officer.	400		
M27	27. Dextran 6% & 0.9% Sodium Chloride IV Injection - 500ml units	To expand plasma volume and maintain blood pressure in the emergency treatment of shock caused by a loss of blood. Also used in the treatment of burns. Dextran should be purchased in a package that includes an administration set.	Give only on RADIO MEDICAL ADVICE from a physician. 500ml infused intravenously at the rate of 20ml per minute, so that the entire amount is given over a period of about 30 minutes. The total amount administered should not exceed 20ml per Kg of body weight during the first 24 hours.	6		
M28	28. Diazepam 5mg/ml Injection (Valium) - 2ml units	(1) For treatment of severe agitation including acute alcohol withdrawal states and convulsions (epilepsy) (2) May be useful for shivering due to generalized hypothermia.	The usual dosage is 2-10 mg (intramuscularly, deep into the muscle), Which may be repeated once in 1-4 hours, depending upon the response and the severity of the condition.	20		
M29	29. Diazepam 5mg tablets(Valium) - 100 tablets	(1) For treatment of common anxiety and tension , and (2) For the management of agitation during alcohol withdrawal.	Varies from 2 to 10mg 2-4 times a day, depending upon the condition being treated.	200		
M30	30. Dimercaprol 100mg/ml Injection - 2ml units	The drug is used as an antidote in poisoning. Indicated in the treatment of arsenic (pesticides), gold and mercury poisonings also in acute lead poisoning, in combination with calcium disodium edetate. Always obtain RADIO MEDICAL ADVICE in the event of such poisoning.	As directed by Doctor or Medical Officer.	6		
M31	31. Doxycycline 100mg Capsules - 50 capsules	Antibacterial agent used in the treatment of urinary infections. It is given in cases of both gonococcal and nongonococcal urethritis after spectinomycin hydrochloride or sulfamethoxazole / trimethoprim has been used.	The usual dose for adults is 200mg on the first day of treatment, given as 100mg every 12hours, followed by a maintenance dose of 100mg daily. In severe infections e.g. chronic infection of the urinary tract, 100mg every 12 hours is recommended	200		



Marshall Islands - Outgoing Up To 24 Persons

Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M32	32. Ephedrine Sulfate 25mg Capsules - 100 capsules	(1) To relieve difficult breathing in asthma (2) to prevent asthmatic attacks in chronic cases (effects appear 30-60 minutes after administration) and (3) to relieve nasal congestion in hayfever and severe head colds. obtain RADIO MEDICAL ADVICE before using this drug	One 25mg capsule four times a day. If ephedrine is used for several days, phenobarbital may be indicated to overcome its stimulant effects (one 30mg phenobarbital tablet up to three times a day)	100		
M33	33. Epinephrine 1mg/ml injection - 1ml units	(1) For acute asthma attacks and for severe allergic reactions to penicillin and other drugs, and to insect bites (2) For cardiac or circulatory failure	By subcutaneous or intramuscular injection, 0.3 ml of 1:1000 solution, every 2 hours as necessary. The usual dosage range is 0.1-0.5 ml in asthma, and up to 1ml in other conditions	20		
M34	34. Methergine 0.2 mg/1ml injection - 1ml units	As Directed	As directed by Doctor or Medical Officer.	10		
M35	35. Erythromycin 250mg tablets - 100 tablets	For a variety of infections of the upper and lower respiratory tract; infections of the mouth, gums, and teeth; infections of the nose, ears, and sinuses. Give this antibiotic only when RADIO MEDICAL ADVICE has been received from a physician, and for some infections resistant to penicillin.	One 250-mg tablet, four times a day (for serious infections, dosage may be increased to two 250mg tablets, four times a day)- to be continued for 48 hours or more after symptoms have subsided.	300		
M36	36. Tetracaine 0.5% Ophth Solution (Pontocaine) - 15ml units	For local anaesthesia of the eye, before removing foreign bodies.	Put 3 drops inside the eyelids, repeat this three times at 2 minute intervals.	2		
M37	37. Triple Antibiotic Pphth Solution - 10ml units	As Directed	As directed by Doctor or Medical Officer.	3		
M38	38. Triple Antibiotic Ophth ointment (Neosporin) 3.5gm	As Directed	As directed by Doctor or Medical Officer.	6		
M39	39. Eye wash Sterile -4oz	For irrigating or flushing the eye to wash away foreign particles, mucous secretions, and fluorescein dye used in diagnosis. Before use, warm the bottle with the solution to body temperature.	As directed by Doctor or Medical Officer.	6		
M40	40. Fluorets Sodium Eye Strips 100 per box	As Directed	As Directed	#		
M41	41. Furosemide tablets, 40mg, (Lasix) 100's.	Furosemide is a potent short-acting diuretic, which causes the production of urine by affecting the kidneys. It is indicated in the treatment of excess body fluid (oedema) associated with congestive heart failure, cirrhosis of the liver, or certain kidney diseases.	The usual oral dosage for adults is 20-80 mg in a single dose in the morning, for 2-4 consecutive days each week, followed by a drug-free period. However, the dosage will vary, depending on the disease and the patients response.	100		
M42	42. Nitro-Quick 0.4mg Sublingual Tablets - 25	As Directed	As directed by Doctor or Medical Officer.	50		
M43	43. Solu-Cortef 100mg Vial Injection (Hydrocortisone) - each	As Directed	As directed by Doctor or Medical Officer.	5		
M44	44. Hydrocortisone 1% Ointment - 1oz	For temporary relief of certain skin disorders, common rashes, inflamed skin and disorders causing itching and discomfort. It may be used for temporary relief of itching, burning and soreness from haemorrhoids (piles).	Apply a thin film to the affected skin 2-4 times a day. Apply sparingly and with gentle rubbing. Clean the skin before each use	6		
M45	45. Ichtammol 10% Ointment - 1oz	As Directed	As directed by Doctor or Medical Officer.	6		
M46	46. Insect repellent pump - 2oz	External use only. To repel mosquitos, chiggers, flies, and other biting insects. It can be used on clothing or the skin, protecting for up to 8 hours.	As directed by Doctor or Medical Officer.	12		
M47	47. Iodine Tincture 2% Mild - 1oz	As Directed	As directed by Doctor or Medical Officer.	10		
M48	Xylocaine 1% 10mg/ml Injection (Lidocaine) - 2ml units	As Directed	As directed by Doctor or Medical Officer.	12		
M49	49. Lindane lotion (Kwell) - 2oz	As Directed	As directed by Doctor or Medical Officer.	12		



Marshall Islands - Outgoing Up To 24 Persons

Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M50	50. Milk of Magnesia - 12oz	As Directed	As directed by Doctor or Medical Officer.	8		
M51	51. Metronidazole 250mg tablets (Flagyl) - 100 tablets	(1) For treatment of salpingitis and other diseases of women and (2) for the treatment of amoebic dysentery.	(1) for diseases of women, usually 1 tablet every 8 hours for 2 weeks. (2) For the dosage in amoebic dysentery, get RADIO MEDICAL ADVICE	500		
M52	52. Miconazole Vaginal Cream (micatin) - 45gm	As Directed	As directed by Doctor or Medical Officer.	5		
M53	53. Miconazole Vaginal Suppositories 7 W/Applicator - each	As Directed	As directed by Doctor or Medical Officer.	3		
M54	54. Mineral oil Heavy - 16oz	Mineral oil (liquid petrolatum) is used as a skin emollient or protective agent. It may also be used to help remove ointments or creams from treated skin areas. Its administration by mouth may be indicated after the ingestion of certain poisons, including alkalis.	Apply liberally over areas of the skin.	1		
M55	55. Morphine sulphate injection, 10mg/ml, 1-ml units	For severe pain not relieved by other analgesics. OBTAIN RADIO MEDICAL ADVICE prior to use. Discontinue as soon as the pain can be relieved by other drugs that can be given orally and are less addictive.	For the relief of severe pain following injuries or burns and severe pain of sudden origin in the abdomen or chest, give 10-15mg intramuscularly. If the pain is unrelieved, or if it recurs soon after the first dose, a second dose of 10 mg intramuscularly, may be given 1 hour or more later. Give further doses if necessary, at intervals of at least 4 hours not sooner.	20		
M56	56. Naloxone Hydrochloride 0.4 mg/ml Injection (Narcan) - 1ml units	For the emergency treatment of respiratory depression resulting from the administration of narcotics, such as morphine. Also indicated for the detection of suspected acute narcotic over-dose.	May be given by subcutaneous, intramuscular, or intravenous injection. The intravenous route of administration is recommended only for emergency situations. the usual dose is 0.4mg, repeated every 2 or 3 minutes until a favourable response is achieved. If no improvement is noted after 2 or 3 doses, another cause of the depression should be suspected.	6		
M57	57. Triple Antibiotic Ointment (Neosporin) - 1oz	As Directed	As directed by Doctor or Medical Officer.	20		
M58	Electrolyte Tablets - 100 tablets	As Directed	As directed by Doctor or Medical Officer.	100		
M59	59. Oxygen Refil Size "D" each	As Directed	As directed by Doctor or Medical Officer.	2		
M60	60. Acetaminophen 500mg tablets (tylenol) - 100 tablets	As Directed	As directed by Doctor or Medical Officer.	300		
M61	61. Petrolatum Ointment - 1oz	As a bland and neutral protective dressing. Apply to minor burns, or dry skin.	As directed by Doctor or Medical Officer.	12		



Marshall Islands - Outgoing Up To 24 Persons

Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M62	62. Phenobarbital 30mg Tablets (Tylenol) - 100 tablets	(1) For insomnia (2) for relatively prolonged sedation, as in some anxiety and tension states and in hypertension and (3) for treatment of epilepsy.	To induce sleep, 90mg at bedtime. As a sedative, one 30mg tablet, 1-4 times daily. In epilepsy, the dose should be adjusted to the individual. consult the doctor in each case. The average range for an adult is 50-120mg daily, although much higher doses may be required for short periods of time	300		
M63	63. Penicillin VK 250mg Tablets - 100 tablets	As Directed	As directed by Doctor or Medical Officer.	300		
M64	64. Pilocarpine 2% Opth Solution (Isopto Carpine) - 15ml units	For the management of glaucoma. Pilocarpine hydrochloride eye-drops constrict the pupil. It is anticipated that this medicament will be needed aboard ship only for patients who have run out of their own supply.	For glaucoma, 2 drops in the eye every 6 hours daily, or as prescribed	2		
M65	65. Potassium Permanganate Crystal - 500g	Diluted un water (1g of crystals to 0.5-1 litre of water.) This substance is used for the disinfection of the skin and mucous membrabes and for the treatment of some skin diseases.	(1g of crystals to 0.5-1 litre of water	1		
M66	66. Probenecid 500mg Tablets (Benemid) - 100 tablets	As an adjunct to intensive therapy with benzylpenicillin, with ampicillin, and a number of other penicillins, for prolonging the effective blood concentration of these drugs	Generally one 500mg tablet four times a day, as long as penicillin is taken.	200		
M67	67. Ampicillin 500mg Injection - each	As Directed	As directed by Doctor or Medical Officer.	20		
M68	68. Proguanil 100mg (Pauludrine) - 100	For prevention of malaria, given either alone or in association with chloroquine. When a ship is in known malarial areas, obtain medical advice on the prophylactic measures effective in that areas.	As directed by Doctor or Medical Officer.	100		
M69	69. Pinworm Caplets (Pyrantel) - 24	For the treatment of (1) pinworm infections and (2) roundworm infections in adults and children	Orally, in a single dose of 10mg per kg of body weight (2 tablets fro a patient weighing 50kg). The highest single dose should not exceed 4 tablets (1g). May be administered with out regard to ingestion of food or time of day. May be taken with either milk or fruit juice	48		
M70	70. Quinine Sulfate 325mg capsules - 100 capsules	In the treatment of acute attacks of malaria contracted in certain areas. Quinine sulfate is not administered to prevent or suppress infection.	For acute malaria attacks, give 2 tablets every 8 hours, until the patient sees a doctor. If the treatment has to be continued for more than 10 days, get RADIO MEDICAL ADVICE. This medicine should be taken after meals to lessen the possibility of stomache upsets.	200		
M71	71. Quinine Dihydrochloride Injection 300mg/ml 2ml units	In the treatment of acute attacks of malaria when the patient is unable to tolerate oral medication because of repeated vomiting or impairment of consciousness.	For acute malaria attacks in adult patients unable to take oral drugs, give 600mg intramuscularly (in the upper outer quadrant of the buttock) every 8 hours until the patient can take oral medication. At that time stop the injections and give quinine sulfate tablets.	20		



Marshall Islands - Outgoing Up To 24 Persons						
Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M72	72. Albuterol 17g Inhalation (Proventil) - 200 metered doses	As Directed	As directed by Doctor or Medical Officer.	2		
M73	73. Sodium Chloride 0.9% IV Injection - 1000ml units	Intravenously (1) to replace water, chloride, and sodium lost in extensive vomiting or diarrhoea, (2) in dehydration due to excessive sweating and (3) in mild haemorrhage. OBTAIN RADIO MEDICAL ADVICE before administering.	Depends on patients condition. Generally, 1500-3000 ml may be given over 24 hours. Do not give intravenous solutions at a faster rate than 500ml per hour, except on RADIO MEDICAL ADVICE	6		
M74	74. IV Solution set vented W/Injection site 15 drop/ml each	As Directed	As directed by Doctor or Medical Officer.	6		
M75	75. IV Solution set 15 Drops/ml Macro drip - each	As Directed	As directed by Doctor or Medical Officer.	6		
M76	76. Thermotabs (enteric coated salt tablets) - 100 tablets	As Directed	As directed by Doctor or Medical Officer.	200		
M77	77. Trobicin Sterile Powder Injection (Spectinomycin)	As Directed	As directed by Doctor or Medical Officer.	20		
M78	78. Sulfamethoxazole/Trimethoprim 400mg/80mg tabs (Bactrim) - 100 tablets	For the treatment of urinary infections, after obtaining RADIO MEDICAL ADVICE	1-2 tablets every 12 hours, for 7 days. Give each dose with full glass of water.	200		
M79	79. Baby Powder J & J (Talc) - 4oz	As as skin lubricant and protector, and as a dusting powder for such skin irritations as chafing and prickly heat	As directed by Doctor or Medical Officer.	6		
M80	80. Tetanus immune Globulin (human) Prefilled syringe - 250 units	To provide passive immunity to tetanus in a patient who is not protected through vaccination with adsorbed tetanus toxoid and has sustained a potentially contaminated wound.	For passive immunization, 250 units by intramuscular injection only.	5		
M81	81. Tetanus Toxoid Prefilled syringe Injection - 0.5ml units	For active immunization against tetanus	For primary immunization of adults and children over 6 years of age. 3 doses are required (see package insert for amount of each dose-usually 0.5 or 10.0ml). The toxoid is injected intramuscularly into the lateral aspect of the upper arm. The second dose is given 4-6 weeks after the first, and the third dose is 6 months to 1 year after the second thereafter, the recommended dose is administered every 5 years.	10		
M82	82. Antibiotic Otic Solution (Generic Cortisporin) - 10ml units	As Directed	As directed by Doctor or Medical Officer.	10		
M83	83. Tetracycline 250mg capsules (Achromycin) - 100 capsules	For susceptible infections; to be given after receiving RADIO MEDICAL ADVICE. do not give to children under 12 years of age.	Usually, one 250mg capsule every 6 hours. In severe illness, increase to two 250-mg capsules every 6 hours.	300		
M84	84. Water for injection Sterile - 5ml units	To dissolve penicillin in powder form	As directed by Doctor or Medical Officer.	30		
M85	85. Zinc oxide ointment	A nontoxic, protective, water-insoluble mildly astringent, and antiseptic paste employed in a large variety of diseases and irritations of the skin.	As directed by Doctor or Medical Officer.	12		
M86	Emesis Basin - Each	As Directed	As Directed	1		
M87	87. Wash Basin Plastic - Large	As Directed	As Directed	1		
M88	88. Bed pan Fracture - each	As Directed	As Directed	1		
M89	89. Cane Aluminum Adjustable #A756 each	As Directed	As Directed	1		
M90	90. Crutches Aluminum Complete - adult regular	As Directed	As Directed	1		
M91	91. Eye Cup Glass each	As Directed	As Directed	2		
M92	92. Funnel Stainless steel - 6oz	As Directed	As Directed	1		
M93	93. Cylinder Glass Double Scale - 50ml	As Directed	As Directed	1		



Marshall Islands - Outgoing Up To 24 Persons

Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M95	95. Heating Pad Cara Dry #50 - each	As Directed	As Directed	1		
M96	96. Hot water bottle / icebag 2 quart each	As Directed	As Directed	1		
M97	97. Strokes litter basket - each	As Directed	As Directed	1		
M98	98. Magnifying glass 2-1/2" - each	As Directed	As Directed	1		
M99	99. Stretcher neil robertson - each	As Directed	As Directed	1		
M100	100. Oxygen unit (complete) W/Case fro "D" size tank - each	As Directed	As Directed	1		
M101	101. Oxygen Cylinder Size "D" empty - each	As Directed	As Directed	1		
M102	102. Refrigerator Walnut colour - Each	As Directed	As Directed	1		
M103	103. Restraints Wrist/Ankle padded pair - each	As Directed	As Directed	1		
M104	104. Straight Jacket	As Directed	As Directed	1		
M105	105. Bedrails Home style #32-GF6408 - 1pair	As Directed	As Directed	1		
M106	106. Resuscitator bag adult disp. w/Mask & tubing	As Directed	As Directed	1		
M107	107. Inflatable ring rubber	As Directed	As Directed	1		
M108	108. Scale doctor plus 0-300lb	As Directed	As Directed	1		
M109	109. Sphygmomanometer aneroid #115	As Directed	As Directed	1		
M110	110. Splint inflatable kit - 4 per kit	As Directed	As Directed	1		
M111	111. Finger Splint padded assorted sizes	As Directed	As Directed	2		
M112	112. Sterilizer Dri-Clave 15-1/2x10-1/2x9-1/2 115 volt	As Directed	As Directed	1		
M113	113. Stethoscope Black - 22"	As Directed	As Directed	1		
M114	114. Thermometer Bath flat wood #50-1537	As Directed	As Directed	2		
M115	115. Instrument Tray Stainless steel - 8-1/2 x 3x1-1/2	As Directed	As Directed	1		
M116	116. Urinal male disposable	As Directed	As Directed	2		
M117	117. Waste recepticle step on 32 Qt white #58-8286	As Directed	As Directed	1		
M118	118. International medical guide for ships	As Directed	As Directed	1		
M119	119. Medical first aid/dangerous goods	As Directed	As Directed	1		
M120	120. International Health regulations	As Directed	As Directed	1		
M121	121. Vaccination Certificate Manual (WHO)	As Directed	As Directed	1		
M122	122. Airway Kit Nasopharyngeal - 5 sizes w/case	As Directed	As Directed	1		
M123	123. Scalpel Blades #10	As Directed	As Directed	6		
M124	124. Scalpel Blades #15	As Directed	As Directed	6		
M125	125. Forceps Dressing Bayonet - shaped 7"	As Directed	As Directed	1		
M126	126. Forceps Halsted Mosquito Curved #20 - 2673	As Directed	As Directed	2		
M127	127. Forceps Kelly Straight - 5-1/2"	As Directed	As Directed	2		
M128	128. Forceps Kelly Curved -5-1/2"	As Directed	As Directed	2		
M129	129. Forceps Splinter -3-1/2"	As Directed	As Directed	1		
M130	130. Forceps Tissue 1x2 teeth -4-1/2"	As Directed	As Directed	1		
M131	131. Scalpel Handle #3	As Directed	As Directed	2		
M132	132. Scissors Bandage -7-1/2"	As Directed	As Directed	3		
M133	133. Scissors Operating Straight Sharp/Sharp-5-1/2"	As Directed	As Directed	2		
M134	134. Hammer Taylor Type Reflex #15-1305-1	As Directed	As Directed	1		
M135	135. Needle Holder Mayo-Hager 5-1/2"	As Directed	As Directed	2		
M136	136. Tape Micropore Paper 2" x 10yd	As Directed	As Directed	4		
M137	137. Tape Micropore Paper 1" x 10 yd	As Directed	As Directed	4		



Marshall Islands - Outgoing Up To 24 Persons						
Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M138	138. IV Butterfly Needle W/Tubing 19g x 3/4"-12"	As Directed	As Directed	6		
M139.	139. Coton Tipped Applicators 6	As Directed	As Directed	200		
M140	140. Elastic Bandage 4" x 4.5 yd	As Directed	As Directed	12		
M141	141. Elastic Bandage 3" x 4.5 yd	As Directed	As Directed	12		
M142	142. Elastic Bandage 2" x 4.5 yd	As Directed	As Directed	12		
M143	143. Elastic Bandage Cotton 2	As Directed	As Directed	12		
M144	144. Flexilite Conforming Gauze Bandage 2" x 4-1/2"	As Directed	As Directed	200		
M145	145. Flexilite Conforming Gauze Bandage 6" x4-1/2"	As Directed	As Directed	40		
M146	146. Flexilite Conforming Gauze Bandage 4" x4-1/2"	As Directed	As Directed	60		
M147	147. Gauze Telfa	As Directed	As Directed	3000		
M148	148. Band Aid Adhesive Surgical Dressing 8" x 6"	As Directed	As Directed	20		
M149	149. Triangular Bandage	As Directed	As Directed	4		
M150	150. Surgitube #2 7/8" x 5 yd	As Directed	As Directed	10		
M151	151. Bandage Spray - 3oz	As Directed	As Directed	4		
M152	152. Vaseline Dressing 3" x 18"	As Directed	As Directed	5		
M153	153. Vaseline Dressing 3" x9"	As Directed	As Directed	10		
M154	154. Vaseline Dressing 6" x 36"	As Directed	As Directed	5		
M155	155. Bandage Compress 4" (1 per box)	As Directed	As Directed	20		
M156	156. Bandage Compress 2" (4 per box)	As Directed	As Directed	3		
M157	157. Bandage Compress 3" (2 per box)	As Directed	As Directed	10		
M158	158. Bottle Plastic - 2oz	As Directed	As Directed	50		
M159	159. Catheter Urethral - 12Fr	As Directed	As Directed	1		
M160	160. Catheter Urethral - 14 Fr	As Directed	As Directed	1		
M161	161. Catheter Urethral - 6 Fr	As Directed	As Directed	1		
M162	162. Catheter Urethral - 8 Fr	As Directed	As Directed	1		
M163	163. Catheter Urethral - 16 Fr	As Directed	As Directed	1		
M164	164. Catheter Tray Urethral - 14 Fr	As Directed	As Directed	3		
M165	165. Clinical Record Chart	As Directed	As Directed	50		
M166	166. Temperature Chart	As Directed	As Directed	50		
M167	167. Medical Report For Seafarers	As Directed	As Directed	100		
M168	168. Cervical Collar Universal - 3"	As Directed	As Directed	1		
M169	169. Cotton Rolled Sterile - 2oz	As Directed	As Directed	3		
M170	170. Cotton Rolled Sterile - 1/2oz	As Directed	As Directed	20		
M171	171. Cotton Rolled Sterile -4oz	As Directed	As Directed	10		
M172	172. Finger Cots Assorted Sizes Sm.,Med., & Large - 12	As Directed	As Directed	12		
M173	173. Penlight Heavy Duty w/batteries	As Directed	As Directed	1		
M174	174. Surgitube #1 5/8" x 5yd	As Directed	As Directed	5		
M175	175. Gloves Latex Surgical "sterile" size 8	As Directed	As Directed	12		
M176	176. Gown Surgical Disposable	As Directed	As Directed	5		
M177	177. Face mask w/elastic band	As Directed	As Directed	50		
M178	178. Medicine Cups plastic 1oz	As Directed	As Directed	100		
M179	179. Microscope slides 1x3 plain	As Directed	As Directed	100		
M180	180. Body bag	As Directed	As Directed	2		
M181	181. Needle Disposable 25g x 5/8"	As Directed	As Directed	36		
M182	182. Needle Disposable 18g x 1-1/2"	As Directed	As Directed	18		
M183	183. Needle Disposable 21g x 1-1/2"	As Directed	As Directed	12		
M184	184. Hematest Reagent tablets	As Directed	As Directed	100		
M185	185. Surgipad Combine Dressing 8"x 10" sterile	As Directed	As Directed	6		
M186	186. Eye pad large sterile - 12 per box	As Directed	As Directed	12		
M187	187. Gauze pads non-adherent 3"x4" sterile - 100 per box	As Directed	As Directed	100		
M188	188. Safety pins Assorted sizes - 50 per	As Directed	As Directed	100		
M189	189. Brush (Surgeons scrub)	As Directed	As Directed	2		
M190	190. Condoms Lubricated	As Directed	As Directed	50		

**Marshall Islands - Outgoing Up To 24 Persons**

Item Number	Item Name	Treatment Requirements	Recommended Dosage	Minimum Qty Required	On Board Quantity	Order Quantity
M191	191. Sheet waterproof 36x72"	As Directed	As Directed	2		
M192	192. Butterfly Closure Medium - 100 each	As Directed	As Directed	1		
M193	193. Steri Strip 1/2"x4" - 6 per	As Directed	As Directed	12		
M194	194. Sutures Chromie Gut 2-0 27" FS1 Needle #637G	As Directed	As Directed	12		
M195	195. Sutures Silk 3-0 18" FS-2 Needle #684G	As Directed	As Directed	24		
M196	196. Sutures Chromic Gut 3-0 27" FS2 Needle #636G	As Directed	As Directed	12		
M197	197. Sutures Nylon 5-0 18" FS-2 Needle #661G	As Directed	As Directed	12		
M198	198. Suture Removal Kit	As Directed	As Directed	1		
M199	199. Glass Syringe 2 ml	As Directed	As Directed	4		
M200	200. Glass Syringe 20ml	As Directed	As Directed	2		
M201	201. Glass Syringe 5 ml	As Directed	As Directed	4		
M202	202. Syringe & Needle 3 cc 21g x 1-1/2"	As Directed	As Directed	50		
M203	203. Syringe & Needle 5 cc 21g x 1-1/2"	As Directed	As Directed	30		
M204	204. Syringe & Needle 3 cc 25g x 5/8"	As Directed	As Directed	50		
M205	205. Chemstrip 9 -100	As Directed	As Directed	100		
M206	206. Clinitest Tablets/Urine Sugar - 36 tablets per	As Directed	As Directed	108		
M207	207. Kleenex - 250 sheets per box	As Directed	As Directed	6		
M208	208. Thermometer Dual scale Oral	As Directed	As Directed	6		
M209	209. Tourniquet Grafkette adult size	As Directed	As Directed	1		
M210	210. Tongue Depressors Wood 6" Senior	As Directed	As Directed	100		
M211	211. Towels Poly-lined 18"x26"	As Directed	As Directed	24		
M212	212. Towels Cotton 18"x33" - 12 each	As Directed	As Directed	12		
M213	213. Sterilized pouches tower self seal 3.5x8" - 100 per	As Directed	As Directed	100		
M214	214. "Sharps" disposal box	As Directed	As Directed	1		